

EDITORIAL

Non-enhanced cardiac computed tomography—still an open book

Ronak Rajani, MD, MRCP, BM, Damini Dey, PhD,
and Daniel S. Berman, MD, FACC

See related article, pp. 96–103

The assessment of coronary artery calcium (CAC) by ECG-gated non-contrast enhanced computed tomography (NCCT) has achieved widespread clinical acceptance as a measure of coronary atherosclerosis and future cardiac risk. A CAC score of 0 is associated with an excellent prognosis^{1,2} while elevated levels provide incremental prognostic information above and beyond that of conventional cardiovascular risk factors in both Caucasian^{3–5} and multiethnic populations.⁶ Specifically, the CAC score has been consistently shown to be more predictive of cardiac outcomes as well as all-cause mortality than the Framingham risk score.^{7,8} Guidelines have been increasingly favorable regarding the use of CAC scanning and currently the test is commonly used in asymptomatic patients for predicting future coronary events and in guiding the intensity of preventive therapies.

In this issue of *JNC*, Gupta et al report a novel application of gated NCCT obtained with the same study used for CAC assessment. They demonstrate a relationship between areas of hypo-attenuation on NCCT and irreversible defects on SPECT myocardial perfusion imaging. Using a cut-off of value of 27.2 HU, they report a sensitivity and specificity of 97.4% and 98.7% for the detection of non-reversible perfusion defects on post-stress and rest imaging with technetium-99m sestamibi.

Assessing the presence of prior myocardial infarction (MI) by NCCT could add significantly to the clinical value of NCCT. While the CAC score is

strongly predictive of patient outcome, it is not highly predictive of coronary stenosis.⁹ Knowing that a patient with coronary calcification had prior MI could help in the prediction of significant coronary artery disease from the NCCT. Further, since NCCT is increasingly performed as part of SPECT or PET–MPI studies, knowledge of prior MI could add to the information provided by the combined scans.

There are, however, several issues with the report by Gupta et al that suggest that the observations should be taken as preliminary. For example, they used standard rest sestamibi scanning to define non-reversibility of perfusion defects, and throughout the manuscript refer to the non-reversible defects as chronic MI. As mentioned by the authors, it is well known that hibernating myocardium could show non-reversible defects in patients without prior MI,^{10,11} and that the prediction of prior MI is enhanced by the use of nitroglycerin prior to the rest injection with sestamibi imaging. Given this limitation, it would have been preferable for them to use the term “non-reversible defect” rather than MI throughout the manuscript. Further, as pointed out by the authors, a primary limitation of their study was that on standard rest sestamibi was used as the reference standard for non-reversible perfusion defects. It would be expected that hibernating regions would have been incorrectly classified as infarct by their approach. In addition, “visual assessment of perfusion abnormalities by the referring physicians” was used to define the presence and location of the non-reversible defects. Given the known high variability of visual assessment of SPECT–MPI by experts,^{12,13} significant inaccuracy among the various readers in the detection and location of the perfusion defects must have been present, such that repeat expert visual reading or quantitative analysis of the data would have been preferred.

Thus, while this provisional data are interesting and demonstrate an additional potential use of NCCT, further validation is required. It would be important to examine the findings with more specific infarct imaging modalities such as cardiac magnetic resonance imaging (CMR), FDG-PET, or possibly nitroglycerin-augmented Tc-99m agent SPECT¹⁴ or Tl-201 rest-redistribution SPECT prior to clinical use.

From the Department of Imaging, Cedars-Sinai Medical Center, Los Angeles, CA.

Reprint requests: Daniel S. Berman, MD, FACC, Department of Imaging, Cedars-Sinai Medical Center, 8700 Beverly Blvd., Los Angeles, CA 90048; bermand@cshs.org.

J Nucl Cardiol 2011;18:21–3.
1071-3581/\$34.00

Copyright © 2010 The Author(s). This article is published with open access at Springerlink.com

doi:10.1007/s12350-010-9325-0

If validated in additional studies, information regarding MI using NCCT would likely be of clinical use in patients undergoing stress imaging studies. In patients undergoing PET/CT or SPECT/CT, gated NCCT is frequently performed. The finding of hypo-attenuation on NCCT could provide complementary information to the perfusion study, aiding in distinguishing MI from attenuation artifact. Further, hypo-attenuation on NCCT, along with the coronary calcium score, may add predictive value in estimating the likelihood of coronary disease and, thus, improve the overall accuracy of perfusion imaging. Finally, as noted above, the information regarding prior MI could aid in predicting significant CAD in patients undergoing CAC scanning alone.

There has recently been renewed interest in the additional anatomic measurements, aside from the coronary arteries, that can be obtained from NCCT in order to best predict the likelihood of developing cardiac disease. Nasir et al¹⁵ demonstrated that, by applying a validated algorithm,¹⁶ measurements of left ventricular volume on NCCT correlated with end-diastolic volume obtained by CMR. Lin et al¹⁷ reported a relationship between aortic root dimensions on NCCT and comparative measurements from transthoracic echocardiography and Wolak et al¹⁸ have established normal ranges of thoracic aortic measurements for age, gender, and body surface area.

Another application of NCCT that is gaining momentum is the measurement of epicardial fat volume (EFV). Epicardial and thoracic fat are routinely imaged on NCCT and there is now emerging evidence that EFV is an important parameter in cardiovascular risk stratification and appears to provide incremental prognostic information regarding future cardiac events over that provided by CAC scanning alone.^{19,20} In contrast to coronary calcium that provides a snapshot of the historical lifetime build up of atherosclerotic plaque, EFV may represent the current state of atherogenesis activity within the coronary arteries.⁸ Recent data also suggest an intriguing relationship between EFV and myocardial ischemia.^{21,22} Although the mechanism of the relationship of coronary atherosclerosis and epicardial fat remains unclear, initial studies indicate that peri-coronary epicardial fat is associated with atherosclerosis in the coronary arteries.²³

The study of Gupta et al provides further evidence that the gated NCCT scan should not simply be considered a coronary calcium scan. As additional substantiating studies emerge, it is plausible that measurements such as LV size, thoracic aortic measurements, epicardial fat, and possibly assessment of prior MI may well become a routine part of the NCCT. The radiation dose associated with NCCT is falling²⁴⁻²⁶ and the additional clinical imaging parameters that can be extracted from this data are increasing. The chapter on coronary calcium may soon be

considered written, but the pages on NCCT keep turning. The story is far from complete.

Acknowledgments

This study was supported in part by ACC/BCS, the Eisner Foundation, and Diane and Guilford Glazer.

Dr Ronak Rajani is currently an advanced cardiac imaging fellow funded by the American College of Cardiology and the British Cardiovascular Society.

The authors have indicated that they have no financial conflicts of interest.

Open Access

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Raggi P, Callister TQ, Cooil B, He ZX, Lippolis NJ, Russo DJ, et al. Identification of patients at increased risk of first unheralded acute myocardial infarction by electron-beam computed tomography. *Circulation* 2000;101:850-5.
2. Arad Y, Goodman KJ, Roth M, Newstein D, Guerci AD. Coronary calcification, coronary disease risk factors, C-reactive protein, and atherosclerotic cardiovascular disease events: The St. Francis Heart Study. *J Am Coll Cardiol* 2005;46:158-65.
3. LaMonte MJ, FitzGerald SJ, Church TS, Barlow CE, Radford NB, Levine BD, et al. Coronary artery calcium score and coronary heart disease events in a large cohort of asymptomatic men and women. *Am J Epidemiol* 2005;162:421-9.
4. Taylor AJ, Bindeman J, Feuerstein I, Cao F, Brazaitis M, O'Malley PG. Coronary calcium independently predicts incident premature coronary heart disease over measured cardiovascular risk factors: Mean three-year outcomes in the Prospective Army Coronary Calcium (PACC) project. *J Am Coll Cardiol* 2005;46:807-14.
5. Vliegenthart R, Oudkerk M, Hofman A, Oei HH, van Dijk W, van Rooij FJ, et al. Coronary calcification improves cardiovascular risk prediction in the elderly. *Circulation* 2005;112:572-7.
6. Detrano R, Guerci AD, Carr JJ, Bild DE, Burke G, Folsom AR, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med* 2008;358:1336-45.
7. Shaw LJ, Raggi P, Schisterman E, Berman DS, Callister TQ. Prognostic value of cardiac risk factors and coronary artery calcium screening for all-cause mortality. *Radiology* 2003;228:826-33.
8. Budoff MJ, Shaw LJ, Liu ST, Weinstein SR, Mosler TP, Tseng PH, et al. Long-term prognosis associated with coronary calcification: Observations from a registry of 25,253 patients. *J Am Coll Cardiol* 2007;49:1860-70.
9. Haberl R, Becker A, Leber A, Knez A, Becker C, Lang C, et al. Correlation of coronary calcification and angiographically documented stenoses in patients with suspected coronary artery disease: Results of 1,764 patients. *J Am Coll Cardiol* 2001;37:451-7.

10. Brunken R, Schwaiger M, Grover-McKay M, Phelps ME, Tillisch J, Schelbert HR. Positron emission tomography detects tissue metabolic activity in myocardial segments with persistent thallium perfusion defects. *J Am Coll Cardiol* 1987;10:557-67.
11. Brunken RC, Kottou S, Nienaber CA, Schwaiger M, Ratib OM, Phelps ME, et al. PET detection of viable tissue in myocardial segments with persistent defects at Tl-201 SPECT. *Radiology* 1989;172:65-73.
12. Cerqueira MD, Nguyen P, Staehr P, Underwood SR, Iskandrian AE. Effects of age, gender, obesity, and diabetes on the efficacy and safety of the selective A2A agonist regadenoson versus adenosine in myocardial perfusion imaging integrated ADVANCE-MPI trial results. *JACC Cardiovasc Imaging* 2008;1:307-16.
13. Berman DS, Kang X, Gransar H, Gerlach J, Friedman JD, Hayes SW, et al. Quantitative assessment of myocardial perfusion abnormality on SPECT myocardial perfusion imaging is more reproducible than expert visual analysis. *J Nucl Cardiol* 2009;16:45-53.
14. Galli M, Marcassa C, Imparato A, Campini R, Orrego PS, Giannuzzi P. Effects of nitroglycerin by technetium-99m sestamibi tomoscintigraphy on resting regional myocardial hypoperfusion in stable patients with healed myocardial infarction. *Am J Cardiol* 1994;74:843-8.
15. Nasir K, Katz R, Mao S, Takasu J, Bomma C, Lima JA, et al. Comparison of left ventricular size by computed tomography with magnetic resonance imaging measures of left ventricle mass and volumes: The multi-ethnic study of atherosclerosis. *J Cardiovasc Comput Tomogr* 2008;2:141-8.
16. Mao S, Budoff MJ, Oudiz RJ, Bakhsheshi H, Wang S, Brundage BH. A simple single slice method for measurement of left and right ventricular enlargement by electron beam tomography. *Int J Card Imaging* 2000;16:383-90.
17. Lin FY, Devereux RB, Roman MJ, Meng J, Jow VM, Jacobs A, et al. Assessment of the thoracic aorta by multidetector computed tomography: Age- and sex-specific reference values in adults without evident cardiovascular disease. *J Cardiovasc Comput Tomogr* 2008;2:298-308.
18. Wolak A, Gransar H, Thomson LE, Friedman JD, Hachamovitch R, Gutstein A, et al. Aortic size assessment by noncontrast cardiac computed tomography: Normal limits by age, gender, and body surface area. *JACC Cardiovasc Imaging* 2008;1:200-9.
19. Mahabadi AA, Massaro JM, Rosito GA, Levy D, Murabito JM, Wolf PA, et al. Association of pericardial fat, intrathoracic fat, and visceral abdominal fat with cardiovascular disease burden: The Framingham Heart Study. *Eur Heart J* 2009;30:850-6.
20. Cheng VY, Dey D, Tamarappoo B, Nakazato R, Gransar H, Miranda-Peats R, et al. Pericardial fat burden on ECG-gated noncontrast CT in asymptomatic patients who subsequently experience adverse cardiovascular events. *JACC Cardiovasc Imaging* 2010;3:352-60.
21. Janik M, Hartlage G, Alexopoulos N, Mirzoyev Z, McLean DS, Arepalli CD, et al. Epicardial adipose tissue volume and coronary artery calcium to predict myocardial ischemia on positron emission tomography-computed tomography studies. *J Nucl Cardiol* 2010;17:841-7.
22. Tamarappoo B, Dey D, Shmilovich H, Nakazato R, Gransar H, Cheng V et al. Increased pericardial fat volume measured from noncontrast CT predicts myocardial ischemia by SPECT. *JACC Cardiovasc Imaging* 2010;3:1104-12.
23. Mahabadi AA, Reinsch N, Lehmann N, Altenbernd J, Kalsch H, Seibel RM, et al. Association of pericoronary fat volume with atherosclerotic plaque burden in the underlying coronary artery: A segment analysis. *Atherosclerosis* 2010;211:195-9.
24. Nakazato R, Dey D, Gutstein A, Le Meunier L, Cheng VY, Pimentel R, et al. Coronary artery calcium scoring using a reduced tube voltage and radiation dose protocol with dual-source computed tomography. *J Cardiovasc Comput Tomogr* 2009;3:394-400.
25. Jakobs TF, Wintersperger BJ, Herzog P, Flohr T, Suess C, Knez A, et al. Ultra-low-dose coronary artery calcium screening using multi-slice CT with retrospective ECG gating. *Eur Radiol* 2003;13:1923-30.
26. Stolzmann P, Leschka S, Betschart T, Desbiolles L, Flohr TG, Marincek B, et al. Radiation dose values for various coronary calcium scoring protocols in dual-source CT. *Int J Cardiovasc Imaging* 2009;25:443-51.